

Science Olympiad

*Eastside Invitational
Stevenson High School
2017*

Chemistry Lab

*Part 1 – Written Exam
Part 2 – Lab Component*

Read the directions below. Do not start the test until you have been instructed to do so.

- You may write on this test.
- You may separate this test into multiple sections.
- Only answers recorded on the scantron will be graded.
- You will find reference pages at the beginning of the test that you may refer to at any point during the test or lab.
- All pages of this test must be stapled together and returned at the end of the testing period.

Team Number: _____

School/Team: _____

Student Names: _____

1

18

| | | | | | | | | | | | | | | | | | |
|---------------------------|---------------------------|---------------------------|---------------------------|---------------------------|---------------------------|---------------------------|---------------------------|---------------------------|---------------------------|---------------------------|---------------------------|---------------------------|---------------------------|---------------------------|---------------------------|---------------------------|---------------------------|
| 1 H 1.008 | 2 | | | | | | | | | | | 13 | 14 | 15 | 16 | 17 | 2 He 4.0026 |
| 3 Li 6.94 | 4 Be 9.0122 | | | | | | | | | | | 5 B 10.81 | 6 C 12.011 | 7 N 14.007 | 8 O 15.999 | 9 F 18.998 | 10 Ne 20.180 |
| 11 Na 22.990 | 12 Mg 24.305 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10 | 11 | 12 | 13 Al 26.982 | 14 Si 28.085 | 15 P 30.974 | 16 S 32.06 | 17 Cl 35.45 | 18 Ar 39.948 |
| 19 K 39.098 | 20 Ca 40.078 | 21 Sc 44.956 | 22 Ti 47.867 | 23 V 50.942 | 24 Cr 51.996 | 25 Mn 54.938 | 26 Fe 55.845 | 27 Co 58.933 | 28 Ni 58.693 | 29 Cu 63.546 | 30 Zn 65.38 | 31 Ga 69.723 | 32 Ge 72.630 | 33 As 74.922 | 34 Se 78.97 | 35 Br 79.904 | 36 Kr 83.798 |
| 37 Rb 85.468 | 38 Sr 87.62 | 39 Y 88.906 | 40 Zr 91.224 | 41 Nb 92.906 | 42 Mo 95.95 | 43 Tc (98) | 44 Ru 101.07 | 45 Rh 102.91 | 46 Pd 106.42 | 47 Ag 107.87 | 48 Cd 112.41 | 49 In 114.82 | 50 Sn 118.71 | 51 Sb 121.76 | 52 Te 127.60 | 53 I 126.90 | 54 Xe 131.29 |
| 55 Cs 132.91 | 56 Ba 137.33 | 57-71 * | 72 Hf 178.49 | 73 Ta 180.95 | 74 W 183.84 | 75 Re 186.21 | 76 Os 190.23 | 77 Ir 192.22 | 78 Pt 195.08 | 79 Au 196.97 | 80 Hg 200.59 | 81 Tl 204.38 | 82 Pb 207.2 | 83 Bi 208.98 | 84 Po (209) | 85 At (210) | 86 Rn (222) |
| 87 Fr (223) | 88 Ra (226) | 89-103 # | 104 Rf (265) | 105 Db (268) | 106 Sg (271) | 107 Bh (270) | 108 Hs (277) | 109 Mt (276) | 110 Ds (281) | 111 Rg (280) | 112 Cn (285) | 113 Nh (286) | 114 Fl (289) | 115 Mc (289) | 116 Lv (293) | 117 Ts (294) | 118 Og (294) |

* Lanthanide series

| | | | | | | | | | | | | | | |
|---------------------------|---------------------------|---------------------------|---------------------------|--------------------------|---------------------------|---------------------------|---------------------------|---------------------------|---------------------------|---------------------------|---------------------------|---------------------------|---------------------------|---------------------------|
| 57 La 138.91 | 58 Ce 140.12 | 59 Pr 140.91 | 60 Nd 144.24 | 61 Pm (145) | 62 Sm 150.36 | 63 Eu 151.96 | 64 Gd 157.25 | 65 Tb 158.93 | 66 Dy 162.50 | 67 Ho 164.93 | 68 Er 167.26 | 69 Tm 168.93 | 70 Yb 173.05 | 71 Lu 174.97 |
|---------------------------|---------------------------|---------------------------|---------------------------|--------------------------|---------------------------|---------------------------|---------------------------|---------------------------|---------------------------|---------------------------|---------------------------|---------------------------|---------------------------|---------------------------|

Actinide series

| | | | | | | | | | | | | | | |
|--------------------------|---------------------------|---------------------------|--------------------------|--------------------------|--------------------------|--------------------------|--------------------------|--------------------------|--------------------------|--------------------------|---------------------------|---------------------------|---------------------------|---------------------------|
| 89 Ac (227) | 90 Th 232.04 | 91 Pa 231.04 | 92 U 238.03 | 93 Np (237) | 94 Pu (244) | 95 Am (243) | 96 Cm (247) | 97 Bk (247) | 98 Cf (251) | 99 Es (252) | 100 Fm (257) | 101 Md (258) | 102 No (259) | 103 Lr (262) |
|--------------------------|---------------------------|---------------------------|--------------------------|--------------------------|--------------------------|--------------------------|--------------------------|--------------------------|--------------------------|--------------------------|---------------------------|---------------------------|---------------------------|---------------------------|

TABLE 10.3 van der Waals Constants for Gas Molecules

| Substance | a (L ² -atm/mol ²) | b (L/mol) |
|------------------|---|-------------|
| He | 0.0341 | 0.02370 |
| Ne | 0.211 | 0.0171 |
| Ar | 1.34 | 0.0322 |
| Kr | 2.32 | 0.0398 |
| Xe | 4.19 | 0.0510 |
| H ₂ | 0.244 | 0.0266 |
| N ₂ | 1.39 | 0.0391 |
| O ₂ | 1.36 | 0.0318 |
| Cl ₂ | 6.49 | 0.0562 |
| H ₂ O | 5.46 | 0.0305 |
| CH ₄ | 2.25 | 0.0428 |
| CO ₂ | 3.59 | 0.0427 |
| CCl ₄ | 20.4 | 0.1383 |

RELATIVE DENSITY OF WATER

| S. No. | Temperature (°C) | Relative density | S. No. | Temperature (°C) | Relative density |
|--------|------------------|------------------|--------|------------------|------------------|
| 1 | 4 | 1.000000 | 22 | 25 | 0.997074 |
| 2 | 5 | 0.999992 | 23 | 26 | 0.996813 |
| 3 | 6 | 0.999968 | 24 | 27 | 0.996542 |
| 4 | 7 | 0.999930 | 25 | 28 | 0.996262 |
| 5 | 8 | 0.999876 | 26 | 29 | 0.995974 |
| 6 | 9 | 0.999809 | 27 | 30 | 0.995676 |
| 7 | 10 | 0.999728 | 28 | 31 | 0.995369 |
| 8 | 11 | 0.999633 | 29 | 32 | 0.995054 |
| 9 | 12 | 0.999525 | 30 | 33 | 0.994731 |
| 10 | 13 | 0.999404 | 31 | 34 | 0.994399 |
| 11 | 14 | 0.999271 | 32 | 35 | 0.994059 |
| 12 | 15 | 0.999127 | 33 | 36 | 0.993712 |
| 13 | 16 | 0.998970 | 34 | 37 | 0.993357 |
| 14 | 17 | 0.998802 | 35 | 38 | 0.992994 |
| 15 | 18 | 0.998623 | 36 | 39 | 0.992623 |
| 16 | 19 | 0.998433 | 37 | 40 | 0.992246 |
| 17 | 20 | 0.998232 | 38 | 41 | 0.99186 |
| 18 | 21 | 0.998021 | 39 | 42 | 0.99147 |
| 19 | 22 | 0.997799 | 40 | 43 | 0.99107 |
| 20 | 23 | 0.997567 | 41 | 44 | 0.99066 |
| 21 | 24 | 0.997326 | 42 | 45 | 0.99024 |

Table 1.
Specific Heat of Water (J/g·°C)

| | |
|--------|-------|
| Solid | 2.1 |
| Liquid | 4.184 |
| Gas | 1.9 |

Table 2.
Latent Heat of Water/Phase Changes (kJ/mol)

| | |
|----------------------------------|------|
| ΔH_{fusion} | 6.02 |
| $\Delta H_{\text{vaporization}}$ | 40.7 |

VAPOR PRESSURE OF WATER

| T | P | T | P | T | P | T | P |
|------|--------|------|--------|------|--------|------|--------|
| °C | torr | °C | torr | °C | torr | °C | torr |
| 19.1 | 16.581 | 22.1 | 19.948 | 25.1 | 23.897 | 28.1 | 28.514 |
| 19.2 | 16.685 | 22.2 | 20.070 | 25.2 | 24.039 | 28.2 | 28.680 |
| 19.3 | 16.789 | 22.3 | 20.193 | 25.3 | 24.182 | 28.3 | 28.847 |
| 19.4 | 16.894 | 22.4 | 20.316 | 25.4 | 24.326 | 28.4 | 29.015 |
| 19.5 | 16.999 | 22.5 | 20.440 | 25.5 | 24.471 | 28.5 | 29.184 |
| 19.6 | 17.105 | 22.6 | 20.565 | 25.6 | 24.617 | 28.6 | 29.354 |
| 19.7 | 17.212 | 22.7 | 20.690 | 25.7 | 24.764 | 28.7 | 29.525 |
| 19.8 | 17.319 | 22.8 | 20.815 | 25.8 | 24.912 | 28.8 | 29.697 |
| 19.9 | 17.427 | 22.9 | 20.941 | 25.9 | 25.060 | 28.9 | 29.870 |
| 20.0 | 17.535 | 23.0 | 21.068 | 26.0 | 25.209 | 29.0 | 30.043 |
| 20.1 | 17.644 | 23.1 | 21.196 | 26.1 | 25.359 | 29.1 | 30.217 |
| 20.2 | 17.753 | 23.2 | 21.324 | 26.2 | 25.509 | 29.2 | 30.392 |
| 20.3 | 17.863 | 23.3 | 21.453 | 26.3 | 25.660 | 29.3 | 30.568 |
| 20.4 | 17.974 | 23.4 | 21.583 | 26.4 | 25.812 | 29.4 | 30.745 |
| 20.5 | 18.085 | 23.5 | 21.714 | 26.5 | 25.964 | 29.5 | 30.923 |
| 20.6 | 18.197 | 23.6 | 21.845 | 26.6 | 26.117 | 29.6 | 31.102 |
| 20.7 | 18.309 | 23.7 | 21.977 | 26.7 | 26.271 | 29.7 | 31.281 |
| 20.8 | 18.422 | 23.8 | 22.110 | 26.8 | 26.426 | 29.8 | 31.461 |
| 20.9 | 18.536 | 23.9 | 22.243 | 26.9 | 26.582 | 29.9 | 31.642 |
| 21.0 | 18.650 | 24.0 | 22.377 | 27.0 | 26.739 | 30.0 | 31.824 |
| 21.1 | 18.765 | 24.1 | 22.512 | 27.1 | 27.897 | 30.1 | 32.007 |
| 21.2 | 18.880 | 24.2 | 22.648 | 27.2 | 27.055 | 30.2 | 32.191 |
| 21.3 | 18.996 | 24.3 | 22.785 | 27.3 | 27.214 | 30.3 | 32.376 |
| 21.4 | 19.113 | 24.4 | 22.922 | 27.4 | 27.374 | 30.4 | 32.561 |
| 21.5 | 19.231 | 24.5 | 23.060 | 27.5 | 27.535 | 30.5 | 32.747 |
| 21.6 | 19.349 | 24.6 | 23.198 | 27.6 | 27.696 | 30.6 | 32.934 |
| 21.7 | 19.468 | 24.7 | 23.337 | 27.7 | 27.858 | 30.7 | 33.122 |
| 21.8 | 19.587 | 24.8 | 23.476 | 27.8 | 28.021 | 30.8 | 33.312 |
| 21.9 | 19.707 | 24.9 | 23.616 | 27.9 | 28.185 | 30.9 | 33.503 |
| 22.0 | 19.827 | 25.0 | 23.756 | 28.0 | 28.349 | 31.0 | 33.695 |

Science Olympiad – Chemistry Lab


Part 1 – Written Test

Multiple Choice – Select the best answer for each below. Be sure to transfer your answers to the scantron.

- Which of the following statements about gases is not true?
 - Gases can be easily compressed
 - The distance between gas molecules is large
 - Gases can form homogenous mixtures
 - Gases will expand to fill their container
 - All gases are colorless and odorless under standard conditions
- A gas sample at constant temperature has a volume of 250 mL and 3.50 atm. What volume will the sample occupy if the pressure is changed to 1.55 atm?
 - 111 mL
 - 217 mL
 - 565 mL
 - 1.36 L
 - 5.65 L
- A sealed balloon will break if its volume reaches 4.50 L. At 20°C, the volume of the balloon is 3.79 L. At what temperature will the balloon break if the pressure stays the same?
 - 17°C
 - 24°C
 - 26°C
 - 51°C
 - 75°C
- How many moles of a gas occupy 2.67 L at 2.3 atm and 25°C?
 - 2.5×10^{-3} mol
 - 0.030 mol
 - 0.25 mol
 - 3.0 mol
 - 4.0 mol
- Which of the following gases would have the highest average molecular speed at 25°C?
 - O₂
 - N₂
 - CO₂
 - CH₄
 - SF₆
- In the van der Waals equation, the constants “a” and “b” are
 - Used to correct for the finite volume of gas molecules and the attractive forces between gas molecules
 - Equal to each other for any real gas
 - Used to correct for the difference between Celsius and Kelvin
 - Equal to 1 for ideal gases
 - Used to correct for the fact that collisions of gas molecules are not really completely elastic
- The vapor pressure of a liquid will decrease if
 - The volume of the vapor above the liquid is increased
 - The volume of the liquid is decreased
 - The temperature is decreased
 - The surface area of the liquid is decreased
 - A more volatile liquid is added

8. Which of the following statements is/are true?
- Deviations in the behavior of gases from the ideal-gas equation occur because gas molecules occupy a finite volume in a container
 - Deviations in the behavior of gases from the ideal-gas equation occur because attractions between gas molecules exist
 - Deviations in the behavior of gases from the ideal-gas equation decrease with increasing temperature
- I only
 - II only
 - I and II
 - II and III
 - I, II, and III
9. How much heat is required to convert 100 g of water at 40°C to water vapor at 100°C?
- 227 kJ
 - 418 kJ
 - 226 kJ
 - 25.1 kJ
 - 251 kJ
10. A chemist uses a cylinder with a piston and gas inlet valve. Consider the following change: Inject an additional gas through the gas inlet valve. What will be the consequences for the pressure of the gas and for the number of moles of gas present?
- The pressure of the gas will decrease, and the number of moles of gas present will decrease
 - The pressure of the gas will increase, and the number of moles of gas present will increase
 - The pressure of the gas will decrease, and the number of moles of gas present will increase
 - There will be no changes in the pressure of the gas or in the number of moles
 - The number of moles will stay the same, and the pressure of the gas will decrease
11. According to the ideal-gas equation, which of the following statements is true?
- If gases are mixed, the partial pressure of each lowers the partial pressure of the others
 - For Boyle's law to apply, a gas must be kept at constant pressure
 - The volume of a gas is not changed if it is heated from 0°C to 100°C and at the same volume if the pressure is increased from 750 torr to 850 torr
 - The volume of a gas doubles when the centigrade temperature doubles if all other variables are held constant
 - The volume of a gas decreases by a factor of 2 when the pressure is doubled if all other variables are held constant
12. At STP, 20. microliters of O₂ contain 5.4×10^{16} molecules. How many molecules are in 20. microliters of N₂?
- 5.4×10^{15}
 - 1.0×10^{16}
 - 2.7×10^{16}
 - 5.4×10^{16}
13. A blimp is filled with 5000. L of helium at 28.0°C and 99.7 kPa. What is the mass of helium used?
- 797 g
 810. g
 - 879 g
 - 8.57×10^3 g
14. What is the density of nitrogen gas at STP?
- 0.62 g/L
 - 1.14 g/L
 - 1.25 g/L
 - 2.03 g/L

15. Find the volume of methane that will produce 12 L of hydrogen in the reaction below. Assume temperature and pressure remain constant.




- 4.0 L
 - 12 L
 - 24 L
 - 36 L
16. What is the partial pressure of oxygen, in torr, in the atmosphere when the atmospheric pressure is 760.0 torr?

| <i>Components of Air</i> | <i>Mole Fraction</i> |
|--------------------------|----------------------|
| Nitrogen | 0.781 |
| Oxygen | 0.209 |
| Argon | 0.010 |

- 159
 - 430
 - 601
 - 720
 - 760
17. A gas is heated in a sealed container. Which of the following occur?
- Gas pressure rises
 - Gas density decreases
 - The average distance between molecules increases
 - All of the above
18. A _____ ΔH corresponds to a _____ process. Select all that are true.
- Negative; endothermic
 - Negative; exothermic
 - Positive; exothermic
 - Positive; endothermic
 - Zero; exothermic
19. An amount of heat equal to 3500 J is released from a system. In addition, 1500 J of work is done by the system on the surroundings. What is the change in internal energy of the system?
- 1500 J
 - 2000 J
 - 3500 J
 - 5000 J
 - 5000 J
20. ΔH° for the reaction below is -482 kJ. Calculate the heat released when 12.0 g of $\text{CO}(\text{g})$ reacts completely, according to the following chemical equation:
- $$2\text{CO}(\text{g}) + \text{O}_2(\text{g}) \rightarrow 2\text{CO}_2(\text{g})$$
- 2.89×10^3 kJ
 - 206 kJ
 - 103 kJ
 - 65.7 kJ
 - 482 kJ
21. What is the specific heat of iron if 13.5 J is required to raise the temperature of a 10-g sample by 3K?
- 0.45 J/g-K
 - 2.22 J/g-K
 - 4.05 J/g-K
 - 45 J/g-K
 - 405 J/g-K

22. For which of the species in the following chemical reaction is the enthalpy of formation equal to zero?



- a. Co(s)
- b. H₂(g)
- c. PF₃(g)
- d. HCo(PF₃)₄(l)
- e. Both Co(s) and H₂(g)

23. What is the standard heat of combustion of CH₄(g)? Use the following data:

| Standard Heats of Formation | |
|-----------------------------|---------------|
| CH ₄ (g) | -74.8 kJ/mol |
| CO ₂ (g) | -393.5 kJ/mol |
| H ₂ O(l) | -285.8 kJ/mol |

- a. -890.3 kJ/mol
- b. -604.6 kJ/mol
- c. -252.9 kJ/mol
- d. -182.5 kJ/mol

24. 10.0 kJ of heat are added to one kilogram of iron at 10. °C. What is the final temperature of the iron? The specific heat of iron is 0.45 J/g-°C.

- a. 22 °C
- b. 27 °C
- c. 32 °C
- d. 37 °C

25. Ozone can be destroyed through reaction with

- a. An oxygen radical
- b. UV radiation
- c. Nitrogen gas
- d. Both a & b

26. Chlorofluorocarbons contribute to ozone depletion by

- a. Releasing chlorine radicals
- b. Directly reacting with ozone
- c. Releasing fluorine radicals
- d. Inhibiting the ability of ozone to migrate to areas of low concentration

27. Which of the following is not a common contributing reaction to ozone depletion?

- a. $\text{HO} + \text{O}_3 \rightarrow \text{HO}_2 + \text{O}_2$
- b. $\text{NO} + \text{O}_3 \rightarrow \text{NO}_2 + \text{O}_2$
- c. $\text{ClO} + \text{O} \rightarrow \text{Cl} + \text{O}_2$
- d. $\text{PO}_3 + \text{O} \rightarrow \text{PO}_2 + \text{O}_2$

28. CFC stands for

- a. Chlorinated Freon compound
- b. Chlorofluorocarbon
- c. Carbonated fluorine compound
- d. Caustic fluorine carbohydrate
- e. Carbofluoro compound

29. The concentration of which greenhouse gas has increased steadily over the last few decades?

- a. H₂O
- b. CO
- c. CO₂
- d. H₂O₂
- e. O₂

30. Cl atoms formed via photolysis of C-Cl bonds of CFC's in the stratosphere are particularly effective in destroying ozone at these altitudes because
- Cl atoms absorb UV, which generate O atoms to react with O_2 to produce ozone
 - Cl atoms catalytically convert O_3 to O_2
 - Cl atoms stoichiometrically convert O_3 to O_2
 - Cl atoms react with H atoms, which catalyze conversion of O_2 to O_3
 - Cl atoms react with N atoms, which catalyze conversion of O_2 to O_3

Completion – solve the following problems. Report your answers on the scantron form in the space provided. Be sure to include units and significant figures with your final answer.

31. A 4.22 g sample of copper (II) sulfide was added to excess hydrochloric acid, and the resulting hydrogen sulfide gas was collected over water. What volume of gas was collected at 30.5°C when the atmospheric pressure was 749 torr?

32. What is the heat change that takes place when 36.0 g of water at atmospheric pressure cools from 125°C to $40.^\circ\text{C}$?

Science Olympiad – Chemistry Lab

Part 2 - Lab Component

Background Information: In this lab you will combine two solutions that react in a 1:1 ratio in order to form a product. The stoichiometry of the reaction is as follows:



Task Details: Use calorimetry to determine the enthalpy of reaction of the product, “C”, in kJ/mol. Assume that the calorimeter is perfectly insulating, that the specific heat of the solution is $4.184 \text{ J}/(\text{g}\cdot^\circ\text{C})$, and that the density of the solution is 1.00 g/mL . Write your final answer in the box provided at the bottom of this page with the correct sign, significant figures, and units.

Materials List:

- Coffee-cup calorimeter (maximum volume of mL)
- Thermometer
- Solution A (mL max.)
- Solution B (mL max.)
- Graduated Cylinder
- Distilled water

Disposal: All solutions can be disposed of in the sink with copious amounts of water.

Data: Clearly record all measurements below.

Calculations: Clearly show all calculations below.

| |
|----------------------|
| Final Answer: |
|----------------------|