

2019-2020 ABRHS Science Olympiad

# Chemistry Lab

# Preliminary Test

Acids/Bases and Aqueous Solutions

Written by: Yuying Fan ([20fany@abschools.org](mailto:20fany@abschools.org))



Name: \_\_\_\_\_ Score: \_\_\_\_\_

Directions:

- You have **50 minutes** to complete this test. Do NOT open the test until instructed.
- You are allowed **one 8.5" x 11" double-sided cheat sheet** and **one stand-alone calculator of any type** for this test.
- For calculations, a correct answer earns full credit, regardless of the work shown. However, work is encouraged for potential partial credit in case the answer is incorrect.
- All answers should be given with the correct units and significant digits (deviations of  $\pm 1$  are allowed). **0.5 points will be deducted for further deviations.**
- The back of the cover page is a periodic table. A blank page is included for scratch work. You may remove any pages from the test packet for your convenience.

# PERIODIC TABLE OF THE ELEMENTS

1	<b>H</b>											2	<b>He</b>				
	1.008											4.00					
3	4											10					
<b>Li</b>	<b>Be</b>											<b>F</b>	<b>Ne</b>				
6.94	9.01											19.00	20.18				
11	12											17	18				
<b>Na</b>	<b>Mg</b>											<b>Cl</b>	<b>Ar</b>				
22.99	24.30											35.45	39.95				
19	20	21	22	23	24	25	26	27	28	29	30	31	32	33	34	35	36
<b>K</b>	<b>Ca</b>	<b>Sc</b>	<b>Ti</b>	<b>V</b>	<b>Cr</b>	<b>Mn</b>	<b>Fe</b>	<b>Co</b>	<b>Ni</b>	<b>Cu</b>	<b>Zn</b>	<b>Ga</b>	<b>Ge</b>	<b>As</b>	<b>Se</b>	<b>Br</b>	<b>Kr</b>
39.10	40.08	44.96	47.90	50.94	52.00	54.94	55.85	58.93	58.69	63.55	65.39	69.72	72.59	74.92	78.96	79.90	83.80
37	38	39	40	41	42	43	44	45	46	47	48	49	50	51	52	53	54
<b>Rb</b>	<b>Sr</b>	<b>Y</b>	<b>Zr</b>	<b>Nb</b>	<b>Mo</b>	<b>Tc</b>	<b>Ru</b>	<b>Rh</b>	<b>Pd</b>	<b>Ag</b>	<b>Cd</b>	<b>In</b>	<b>Sn</b>	<b>Sb</b>	<b>Te</b>	<b>I</b>	<b>Xe</b>
85.47	87.62	88.91	91.22	92.91	95.94	(98)	101.1	102.91	106.42	107.87	112.41	114.82	118.71	121.75	127.60	126.91	131.29
55	56	57	72	73	74	75	76	77	78	79	80	81	82	83	84	85	86
<b>Cs</b>	<b>Ba</b>	<b>*La</b>	<b>Hf</b>	<b>Ta</b>	<b>W</b>	<b>Re</b>	<b>Os</b>	<b>Ir</b>	<b>Pt</b>	<b>Au</b>	<b>Hg</b>	<b>Tl</b>	<b>Pb</b>	<b>Bi</b>	<b>Po</b>	<b>At</b>	<b>Rn</b>
132.91	137.33	138.91	178.49	180.95	183.85	186.21	190.2	192.2	195.08	196.97	200.59	204.38	207.2	208.98	(209)	(210)	(222)
87	88	89	104	105	106	107	108	109	110	111							
<b>Fr</b>	<b>Ra</b>	<b>†Ac</b>	<b>Rf</b>	<b>Db</b>	<b>Sg</b>	<b>Bh</b>	<b>Hs</b>	<b>Mt</b>	<b>Ds</b>	<b>Rg</b>							
(223)	226.02	227.03	(261)	(262)	(266)	(264)	(277)	(268)	(271)	(272)							
*Lanthanide Series																	
58	59	60	61	62	63	64	65	66	67	68	69	70	71				
<b>Ce</b>	<b>Pr</b>	<b>Nd</b>	<b>Pm</b>	<b>Sm</b>	<b>Eu</b>	<b>Gd</b>	<b>Tb</b>	<b>Dy</b>	<b>Ho</b>	<b>Er</b>	<b>Tm</b>	<b>Yb</b>	<b>Lu</b>				
140.12	140.91	144.24	(145)	150.4	151.97	157.25	158.93	162.50	164.93	167.26	168.93	173.04	174.97				
90	91	92	93	94	95	96	97	98	99	100	101	102	103				
<b>Th</b>	<b>Pa</b>	<b>U</b>	<b>Np</b>	<b>Pu</b>	<b>Am</b>	<b>Cm</b>	<b>Bk</b>	<b>Cf</b>	<b>Es</b>	<b>Fm</b>	<b>Md</b>	<b>No</b>	<b>Lr</b>				
232.04	231.04	238.03	(237)	(244)	(243)	(247)	(247)	(251)	(252)	(257)	(258)	(259)	(262)				
†Actinide Series																	





**Part I: Multiple Choice (2 pts each, 30 pts total)**

1. Under room temperature, the product of concentrations of  $\text{H}^+$  and  $\text{OH}^-$  produced by the auto-ionization of water in a solution is  $10^{-24} \text{ mol}^2 \cdot \text{L}^{-2}$ . In this solution, \_\_\_\_\_ CANNOT possibly exist in a large quantity.
- A.  $\text{SO}_3^{2-}$
  - B.  $\text{NH}_4^+$
  - C.  $\text{NO}_3^-$
  - D.  $\text{HCO}_3^-$
2. When temperature increases, which of the following species does NOT increase in its solubility in water?
- A.  $\text{NaNO}_3$
  - B. KI
  - C. KCl
  - D.  $\text{NH}_3$
3. x mL of a hydrochloric acid solution with pH = 2.5 will completely react with exactly 10x mL of a strong monoprotic base solution. What is the pH of the basic solution?
- A. 9.0
  - B. 9.5
  - C. 10.5
  - D. 11.0
4. There is 10 mL of a strong acid solution with a pH of 1 under room temperature. How to make the pH of the solution become 2?
- A. Add water to dilute the solution to 100 mL
  - B. Add 10 mL of water to dilute the solution
  - C. Add 10 mL of 0.01M NaOH solution
  - D. Add 10 mL of 0.01M hydrochloric acid solution
5. Under room temperature, some crystals precipitated out of a glass of saturated table salt solution after it was exposed to air for a while. This is because \_\_\_\_\_.
- A. the solubility of NaCl decreased
  - B. the mass fraction of solute in the solution decreased
  - C. the solution became unsaturated
  - D. the mass of the solvent decreased

6. Which of the following groups of solids can be separated following the steps “dissolution → filtration → evaporation”?

- A.  $\text{MnO}_2$  and  $\text{NaCl}$
- B.  $\text{CuO}$  and  $\text{CaCO}_3$
- C.  $\text{Fe}$  and  $\text{S}$
- D.  $\text{KMnO}_4$  and  $\text{KClO}_3$

7. Under room temperature, which of the following statements on solution dilution is correct?

- A. After diluting a 1 L 0.1M  $\text{Ba}(\text{OH})_2$  solution to 2 L by adding water, the final pH is 13.
- B. After diluting a hydrochloric acid solution with a pH of 6 to 10 times its original volume by adding water, the final pH is 7.
- C. After diluting an  $\text{H}_2\text{SO}_4$  solution with a pH of 4 to 100 times its original volume by adding water, the  $[\text{H}^+]$  produced by auto-ionization of water in the solution is  $1 \times 10^{-6}$  M.
- D. After diluting a  $\text{NaOH}$  solution with a pH of 8 to 100 times its original volume by adding water, the final pH is 6.

8. 100 mL of 0.02M  $\text{Ba}(\text{OH})_2$  solution and 100 mL of 0.02M  $\text{NaHSO}_4$  solution are mixed under room temperature. Ignoring the change in volume, in the resulting solution \_\_\_\_\_.

- A.  $\text{pH} = 12$
- B.  $\text{pH} = 2$
- C.  $[\text{H}^+]$  due to the auto-ionization of water is equal to  $1.0 \times 10^{-2}$  M
- D. the total concentration of solutes is 0.02M

9. After mixing equal volumes of two  $\text{NaOH}$  solutions ( $\text{pH} = 8$  and  $\text{pH} = 10$ , respectively), the pH of the resulting solution is closest to which of the following?

- A. 8.3
- B. 8.7
- C. 9.3
- D. 9.7

10. Using distilled water, an acetic acid solution and a hydrochloric acid solution with the same pH are diluted to m times and n times the original volume, respectively. If the pH of the two solutions is still the same after dilution, what is the relationship between m and n?

- A.  $m < n$
- B.  $m = n$
- C.  $m > n$
- D.  $m = 2n$

11. After pouring out  $\frac{1}{2}$  of a 10% dilute sulfuric acid solution from a beaker, the remaining solution \_\_\_\_\_.

- A. has a solute mass fraction of 20%
- B. has a halved solute mass fraction, or 5%
- C. has the same solute mass fraction as before, which is 10%
- D. does not match any of the statements above

12. Let  $n$  be Avogadro's number. For 0.2M sodium sulfate, which of the following is correct?

- A. A 1 L solution contains  $0.2n$  sodium ions.
- B. A 1 L solution contains  $0.6n$  sodium ions and sulfate ions in total.
- C. A 3 L solution has a sodium ion concentration of 1.2M.
- D. A 2 L solution contains  $0.6n$  sulfate ions.

13. Which of the following gives an accurate description of the relationships among concentrations of ions in solution?

- A. In solutions of ammonia, KOH and  $\text{Ba}(\text{OH})_2$  of the same pH,  $[\text{NH}_4^+] = [\text{K}^+] = [\text{Ba}^{2+}]$ .
- B. After adding 10 mL of 0.1M  $\text{Na}_2\text{CO}_3$  dropwise into 10 mL of 0.1M hydrochloric acid,  $[\text{Na}^+] > [\text{Cl}^-] > [\text{HCO}_3^-] > [\text{CO}_3^{2-}]$ .
- C. After adding a NaOH solution dropwise into a  $\text{NH}_4\text{HCO}_3$  solution until pH reaches 7,  $[\text{NH}_4^+] + [\text{Na}^+] = [\text{HCO}_3^-] + [\text{CO}_3^{2-}]$ .
- D. A 0.2M monoprotic acid HA solution is mixed with 0.1M NaOH of the same volume.  $2[\text{OH}^-] + [\text{A}^-] = 2[\text{H}^+] + [\text{HA}]$ .

14. Which of the following groups of ions can coexist in solution at large quantities?

- A.  $\text{Fe}^{3+}$ ,  $\text{Na}^+$ ,  $\text{Cl}^-$  and  $\text{I}^-$
- B.  $\text{Cu}^{2+}$ ,  $\text{K}^+$ ,  $\text{NO}_3^-$  and  $\text{S}^{2-}$
- C.  $\text{AlO}_2^-$ ,  $\text{K}^+$ ,  $\text{NO}_3^-$  and  $\text{OH}^-$
- D.  $\text{Ba}^{2+}$ ,  $\text{H}^+$ ,  $\text{Cl}^-$  and  $\text{SO}_4^{2-}$

15. At 95 °C, the concentration of  $\text{H}^+$  in pure water is  $10^{-6} \text{ mol} \cdot \text{L}^{-1}$ . If 0.01 mol of NaOH solid is dissolved in water at 95 °C to make a 1 L solution, then the pH of the resulting solution is \_\_\_\_\_.

- A. 4
- B. 10
- C. 2
- D. 12

**Part II: Open Response (70 pts total)**

1. Refer to the picture on the cover page. The four characters are Mg, Fe, Cu and Zn, and the solution in the test tubes is hydrochloric acid.

a) Match the characters with their identities in order from left to right. (2 pts)

b) Name the gas formed in the test tubes. (1 pt)

2. Complete the table. (1 pt each, 8 pts total)

Acid			$\text{HPO}_4^{2-}$	$\text{CH}_3\text{COOH}$
Conjugate Base	$\text{OH}^-$	$\text{H}_2\text{O}$		
Acid		$[\text{Hg}(\text{H}_2\text{O})_6]^{2+}$	$[\text{Zn}(\text{H}_2\text{O})_4(\text{OH})_2]$	
Conjugate Base	$\text{C}_2\text{H}_5\text{NH}_2$			$\text{HF}$

3. a) Which of the following is/are characteristic(s) of aqueous solutions? (Write all numbers that apply) (2 pts)

① uniform ② is a mixture ③ colorless ④ transparent ⑤ is a compound

b) Give two examples of solutions from NATURE or DAILY LIFE (note: while it may not necessarily apply to you, an average typical person from the street does not see 0.1M  $\text{H}_2\text{SO}_4$  solution or the like in daily life). (2 pts)

c) What are colligative properties? (2 pts)

d) Give two examples of colligative properties. (2 pts)

4. Illustrate the following processes.

a) NaCl dissolves in water. (2 pts)

b) Fluoride ion reacts with boron trifluoride to make tetrafluoroborate. (2 pts)

\* What type of acid-base reaction is this? \_\_\_\_\_ (1 pt)


\* What is the acid? \_\_\_\_\_ (1 pt)

c) Auto-ionization of water. (2 pts)

d) 100 grams of magnesium chloride mixed with 100 grams of water. (The solubility of magnesium chloride in water is 54.3 g/100 mL at 20 °C.) (2 pts)

e) The neutralization reaction between sodium hydroxide and Vitamin C (show structural formula). (2 pts)

5. The following scene represents three weak acids HA (where A = X, Y, or Z) dissolved in water ( $\text{H}_2\text{O}$  is not shown):



a) Rank the acids in order of increasing  $K_a$ . (1 pt)

b) Rank the acids in order of increasing  $\text{p}K_a$ . (1 pt)

c) Rank the conjugate bases in order of increasing  $\text{p}K_b$ . (1 pt)

d) If equimolar amounts of the sodium salts of the acids ( $\text{NaX}$ ,  $\text{NaY}$ , and  $\text{NaZ}$ ) were dissolved in water, which solution would have the highest  $\text{pOH}$ ? Which solution would have the lowest  $\text{pH}$ ? (2 pt)

6. Consider the following five solutions with the same molar concentration:

①  $\text{CH}_3\text{COONa}$     ②  $\text{CH}_3\text{COONH}_4$     ③  $\text{CH}_3\text{COOH}$     ④  $\text{NH}_3 \cdot \text{H}_2\text{O}$     ⑤  $\text{NH}_4\text{Cl}$

a) Rank the solutions in order of descending  $\text{pH}$  (write the numbers, not names; indicate which ones are equal, if any): (2 pts)

b) Rank the solutions in order of descending total concentrations of ions (write the numbers, not names; indicate which ones are equal, if any): (2 pts)

7. A solution is prepared by adding 29.3 g of menthol ( $C_{10}H_{20}O$ ) to 0.0590 kg of chloroform.

a) What is its boiling point? (The boiling point of pure chloroform is  $61.2\text{ }^{\circ}\text{C}$ , and the boiling point constant for chloroform is  $3.85\text{ }^{\circ}\text{C/m}$ .) (2 pts)

b) What is the concentration of menthol expressed in molarity? (The density of menthol is  $0.89\text{ g/mL}$ , and the density of chloroform is  $1.492\text{ g/mL}$  at  $25\text{ }^{\circ}\text{C}$ .) (1 pt)

8. After diluting an  $H_2SO_4$  solution with a pH of 5 to 500 times its original volume, what is the approximate ratio between  $[H^+]$  and  $[SO_4^{2-}]$ ? (3 pts)

9. 100 mL of a 0.1M  $Ba(OH)_2$  solution and 50 mL of a 0.1M  $Na_2CO_3$  solution are mixed together. A student sets up a titration and adds 0.1M HCl into the mixture dropwise while swirling.

a) When the pH reaches 7, the student stops. What is the volume of HCl added? (2 pts)

b) If the student then adds 0.1M acidified  $AgNO_3$  into the solution until all  $Cl^-$  completely precipitates, what is the minimum volume of  $AgNO_3$  solution needed? (2 pts)

10. What is the pH of a 0.300M solution of benzoic acid ( $K_a = 6.46 \times 10^{-5}$ )? (2 pts)

11. How many grams of sodium formate ( $\text{HCOONa}$ ) do you need to add to 400. mL of 1.00M formic acid to form a buffer with  $\text{pH} = 3.500$ ? ( $K_a = 1.77 \times 10^{-4}$ ) (2 pts)

12. When using neutralization titration to determine the content of NaOH in a mixture of NaOH and  $\text{Na}_2\text{CO}_3$  solutions, a student first adds excess barium chloride to completely precipitate out  $\text{Na}_2\text{CO}_3$  as  $\text{BaCO}_3$ , then titrate the solution with standardized hydrochloric acid (using phenolphthalein as the indicator).

a) Why will adding hydrochloric acid into the NaOH solution mixed with  $\text{BaCO}_3$  precipitate not cause the  $\text{BaCO}_3$  to dissolve but allow the student to determine the content of NaOH? (2 pts)

b) How will the color of the indicator change at the endpoint? (1 pt)

c) During the titration, if the endpoint was not reached until the titrant level fell under the stopcock of the buret, the recorded volume of hydrochloric acid used will be \_\_\_\_\_ (too high/accurate/too low). (1 pt)

d) Can the student use methyl orange (pH range = 3.1–4.4) as the indicator instead? \_\_\_\_\_ (Yes/No) (1 pt). If methyl orange is used as the indicator, the content of NaOH calculated based on the results will be \_\_\_\_\_ (too high/accurate/too low) (1 pt).

13. One of the raw materials of the chlor-alkali industry, saturated table salt solution, contains a certain quantity of ammonium ions. During electrolysis, these ammonium ions will form the extremely unstable compound nitrogen trichloride, which can easily cause an explosion.

a) Nitrogen trichloride is readily hydrolyzed. Aside from ammonia gas, its initial products from hydrolysis also include \_\_\_\_\_. (1 pt)

b) To remove the ammonium ions from saturated table salt solution, one method is to introduce chlorine gas under basic conditions to react and form nitrogen gas. The net ionic equation for this reaction is \_\_\_\_\_ (2 pts).

One advantage of choosing this method in industry is \_\_\_\_\_ (1 pt).

c) Excess chlorine gas is removed using  $\text{Na}_2\text{S}_2\text{O}_3$ . In the reaction,  $\text{S}_2\text{O}_3^{2-}$  is oxidized and becomes  $\text{SO}_4^{2-}$ . If there are  $1 \times 10^{-3}$  moles of chlorine gas in excess, then theoretically the  $\text{SO}_4^{2-}$  produced will be \_\_\_\_\_ mol. (3 pts)

d) Formaldehyde is widely used in production and experiments to determine the nitrogen content of a sample. The reaction between formaldehyde and ammonium ion is  $4\text{NH}_4^+ + 6\text{HCHO} \rightarrow (\text{CH}_2)_6\text{N}_4\text{H}^+ + 3\text{H}^+ + 6\text{H}_2\text{O}$ , with  $(\text{CH}_2)_6\text{N}_4\text{H}^+$  being a monoprotic acid.

The procedures are as follows:

① Formic acid is often contained in formaldehyde solutions. Pour  $b$  mL of formaldehyde solution (excess) into an Erlenmeyer flask and add one drop of phenolphthalein. Neutralize the solution with  $c$  mol/L NaOH. The initial reading on the buret is  $V_1$  mL; when the solution appears to be light pink, the reading on the buret  $V_2$  mL.

② Add  $a$  mL of a sample of saturated table salt solution and let the flask stand for several minutes.

③ Add 1–2 more drops of phenolphthalein, and use the same NaOH solution to titrate until light pink shows. The reading on the buret is  $V_3$  mL.

The nitrogen content of the sample of table salt solution is \_\_\_\_\_ mg/L (express in terms of the variables). (3 pts)